

## References

- Anderson, P.J., Rogers, Q.R. and Morris, J.G., 2002. Cats require more dietary phenylalanine or tyrosine for melanin deposition in hair than for maximal growth. *The Journal of nutrition*, 132(7), pp.2037-2042.
- Bosch, G., Beerda, B., Hendriks, W.H., Van der Poel, A.F.B. and Verstegen, M.W.A., 2007. Impact of nutrition on canine behaviour: current status and possible mechanisms. *Nutrition research reviews*, 20(2), pp.180-194.
- Broer, S., 2008. Amino acid transport across mammalian intestinal and renal epithelia. *Physiological reviews*, 88(1), pp.249-286.
- Chipponi, J.X., Bleier, J.C., Santi, M.T. and Rudman, D., 1982. Deficiencies of essential and conditionally essential nutrients. *The American journal of clinical nutrition*, 35(5), pp.1112-1116.
- Johnson, M. L., C. M. Parsons, G. C. Fahey Jr, N. R. Merchen, and C. G. Aldrich. 1998. Effects of species raw material source, ash content, and processing temperatures on amino acid digestibility of animal by-product meals by cecectomized roosters and ideally cannulated dogs. *J. Anim. Sci.* 76:1112-1122. doi:10.2527/1998.7641112x.
- Lambie, J.G., Pezzali, J.G., Richards, T.L., Ellis, J.L., Verbrugghe, A. and Shoveller, A.K., 2024. Phenylalanine requirements using the direct amino acid oxidation technique, and the effects of dietary phenylalanine on food intake, gastric emptying, and macronutrient metabolism in adult cats. *Journal of Animal Science*, 102, p.skae009.
- Mansilla, W.D., Templeman, J.R., Fortener, L. and Shoveller, A.K., 2020. Minimum dietary methionine requirements in Miniature Dachshund, Beagle, and Labrador Retriever adult dogs using the indicator amino acid oxidation technique. *Journal of animal science*, 98(11), p.skaa324.
- Morris, J.G., Yu, S. and Rogers, Q.R., 2002. Red hair in black cats is reversed by addition of tyrosine to the diet. *The journal of Nutrition*, 132(6), pp.1646S-1648S.
- National Research Council. 2006. Nutrient requirements of dogs and cats. 2nd rev. ed. Washington (DC): National Academic Press.
- Pavlova, N.N., King, B., Josselsohn, R.H., Violante, S., Macera, V.L., Vardhana, S.A., Cross, J.R. and Thompson, C.B., 2020. Translation in amino-acid-poor environments is limited by tRNAGln charging. *Elife*, 9, p.e62307.
- Pencharz, P.B. and Ball, R.O., 2003. Different approaches to define individual amino acid requirements. *Annual review of nutrition*, 23(1), pp.101-116.
- Pezzali, J.G., Lambie, J.G., Verbrugghe, A. and Shoveller, A.K., 2024. Minimum methionine requirement in adult cats as determined by indicator amino acid oxidation. *Journal of Animal Science*, 102, p.skad411.
- Reeds PJ (2000) Dispensable and indispensable amino acids for humans. *J Nutr* 130:1835S-1840S
- Sutherland, K.A., Mansilla, W.D., Fortener, L. and Shoveller, A.K., 2020. Lysine requirements in small, medium, and large breed adult dogs using the indicator amino acid oxidation technique. *Translational Animal Science*, 4(3), p.txa082.
- Templeman, J.R., Mansilla, W.D., Fortener, L. and Shoveller, A.K., 2019. Tryptophan requirements in small, medium, and large breed adult dogs using the indicator amino acid oxidation technique. *Journal of animal science*, 97(8), pp.3274-3285.
- Wu, G., Bazer, F.W., Dai, Z., Li, D., Wang, J. and Wu, Z., 2014. Amino acid nutrition in animals: protein synthesis and beyond. *Annu. Rev. Anim. Biosci.*, 2(1), pp.387-417.
- Yu, S., Rogers, Q.R. and Morris, J.G., 2001. Effect of low levels of dietary tyrosine on the hair colour of cats. *Journal of Small Animal Practice*, 42(4), pp.176-180.
- Zello, G.A., Wykes, L.J., Ball, R.O. and Pencharz, P.B., 1995. Recent advances in methods of assessing dietary amino acid requirements for adult humans. *The Journal of nutrition*, 125(12), pp.2907-2915.
- Zhang, S., de Vries, S. and Gerrits, W.J., 2024. Quantifying the effects of dietary fibres on protein digestibility in pigs-A review. *Animal Feed Science and Technology*, 308, p.115864.